

# IMMUNIZATION AWARENESS

## AUGUST 2024



**SIHO** INSURANCE SERVICES

The CDC has several immunizations that are currently recommended for adults over the age of 18. Per the Indiana State Department of Health, the immunizations that you need qualify for are different based on several different factors such as:

- Your immunization history
- Age and lifestyle, and any existing health conditions (diabetes, autoimmune disorders, etc.)
- Certain occupations require additional vaccines
- International travel

VACCINE	19-26 YEARS	27-49 YEARS	50-64 YEARS	60+ YEARS
<b>COVID-19</b>	1 or more doses of updated (2023–2024 Formula) vaccine			
<b>Influenza inactivated (IIV4)</b> or <b>Influenza recombinant (RIV4)</b>	1 dose annually			
<b>or</b>	<b>or</b>			
<b>Influenza live attenuated (LAIV4)</b>	1 dose annually			
<b>Respiratory Syncytial Virus (RSV)</b>	Seasonal administration during pregnancy.			≥ 60 years
<b>Tetanus, diphtheria, pertussis (Tdap or Td)</b>	1 dose Tdap each pregnancy; 1 dose Td/Tdap for wound management			
	1 dose Tdap, then Td or Tdap booster every 10 years			
<b>Measles, mumps, rubella (MMR)</b>	1 or 2 doses depending on indication (if born in 1957 or later)			
<b>Varicella (VAR)</b>	2 doses (if born in 1980 or later)		2 doses	
<b>Zoster recombinant (RZV)</b>	2 doses for immunocompromising conditions		2 doses	
<b>Human papillomavirus (HPV)</b>	2 or 3 doses depending on age at initial vaccination or condition	27 through 45 years		
<b>Pneumococcal (PCV15, PCV20, PPSV23)</b>	1 dose PCV15 followed by PPSV23 OR 1 dose PCV20			1 dose PCV15 followed by PPSV23 OR 1 dose PCV20
<b>Hepatitis A (HepA)</b>	2 or 3 doses depending on vaccine			
<b>Hepatitis B (HepB)</b>	2, 3, or 4 doses depending on vaccine or condition			
<b>Meningococcal A, C, W, Y (MenACWY)</b>	1 or 2 doses depending on indication			
<b>Meningococcal B (MenB)</b>	19 through 23 years	2 or 3 doses depending on vaccine and indication		
<b>Haemophilus influenzae type b (Hib)</b>	1 or 3 doses depending on indication			
<b>Mpox</b>				

Recommended vaccination for adults who meet age requirement, lack documentation of vaccination, or lack evidence of immunity.

Recommended vaccination for adults with an additional risk factor or another indication.

Recommended vaccination based on shared clinical decision-making.

No recommendation/Not applicable